

## Pointwise error estimates for a singularly perturbed time-dependent semilinear reaction-diffusion problem\*

NATALIA KOPTOVA AND SIMONA BLANCA SAVESCU†

*Mathematics and Statistics Department, University of Limerick,  
Limerick, Ireland.*

An initial-boundary-value problem for a semilinear reaction-diffusion equation is considered. Its diffusion parameter  $\varepsilon^2$  is arbitrarily small, which induces initial and boundary layers. It is shown that the conventional implicit method might produce incorrect computed solutions on uniform meshes. Therefore we propose a stabilized method that yields a unique qualitatively-correct solution on any mesh. Constructing discrete upper and lower solutions, we prove existence and investigate the accuracy of discrete solutions on layer-adapted meshes of Bakhvalov and Shishkin types. It is established that the two considered methods enjoy second-order convergence (with, in the case of the Shishkin mesh, a logarithmic factor) in the discrete maximum norm, uniformly in  $\varepsilon$  for  $\varepsilon \leq C(N^{-1} + M^{-1/2})$ , where  $N$  and  $M$  are the numbers of mesh intervals in the space and time directions, respectively. Numerical results are presented that support the theoretical conclusions.

*Keywords:* semilinear reaction-diffusion, singular perturbation, maximum norm error estimate, Bakhvalov mesh, Shishkin mesh, second order, upper and lower solutions.

### 1. Introduction

Consider the singularly perturbed semilinear reaction-diffusion equation

$$\mathcal{T}u \equiv \varepsilon^2[u_t - u_{xx}] + f(x, t, u) = 0 \quad \text{for } (x, t) \in (0, 1) \times (0, T], \quad (1.1a)$$

subject to the boundary and initial conditions

$$u(0, t) = g_0(t), \quad u(1, t) = g_1(t), \quad t \in [0, T], \quad (1.1b)$$

$$u(x, 0) = \varphi(x), \quad x \in [0, 1]. \quad (1.1c)$$

Here  $\varepsilon$  is a small positive parameter, and the functions  $f$ ,  $g_0$ ,  $g_1$  and  $\varphi$  are sufficiently smooth; furthermore, at the corners  $(0, 0)$  and  $(1, 0)$  of our domain we assume the standard compatibility conditions  $g_0(0) = \varphi(0)$  and  $g_1(0) = \varphi(1)$ .

In the numerical analysis literature it is often assumed that  $f_u(x, t, u) > 0$  for all  $(x, t, u) \in [0, 1] \times [0, T] \times \mathbb{R}$ . This global condition is nevertheless rather restrictive. E.g., mathematical models of biological and chemical processes frequently involve problems related to (1.1) with  $f(x, t, u)$  that is *non-monotone* with respect to  $u$ ; see, e.g., (Murray, 1993, §14.7), (Grindrod, 1991, §2.3). Hence we drop the assumption that  $f_u > 0$  and consider problem (1.1) under weaker assumptions, described in §2, that intrinsically arise from the asymptotic analysis of this problem.

\*This publication has emanated from research conducted with the financial support of Science Foundation Ireland under the Research Frontiers Programme 2008; Grant 08/RFP/MTH1536.

†Email: Natalia.Kopteva@ul.ie; Simona.Savescu@ul.ie

The reduced problem of (1.1) is defined by formally setting  $\varepsilon = 0$  in (1.1a), i.e.

$$f(x, t, u_0(x, t)) = 0 \quad \text{for } (x, t) \in (0, 1) \times (0, T). \quad (1.2)$$

As  $f_u$  is not necessarily positive, this equation might have multiple solutions, and any solution  $u_0$  of (1.2) does not in general satisfy the boundary and initial conditions in (1.1b) and (1.1c). Similarly, the steady-state version of (1.1) might have multiple solutions. In contrast, the initial-boundary-value problem (1.1) always has at most one solution; see Proposition 1.1 below. Therefore, if problem (1.1) is solved numerically, it is desirable that the computed solution enjoys a similar property.

We discretize (1.1) on a tensor-product mesh  $\{(x_i, t_j)\}$  in  $[0, 1] \times [0, T]$ , where  $0 = x_0 < x_1 < \dots < x_N = 1$  and  $0 = t_0 < t_1 < \dots < t_M = T$ , and we use the notation  $h_i := x_i - x_{i-1}$  and  $k_j = t_j - t_{j-1}$  for the local mesh sizes. One standard implicit discretization of (1.1) is given by

$$\mathcal{J}^h U_{ij} := \varepsilon^2 [\delta_t - \delta_x^2] U_{ij} + f(x_i, t_j, U_{ij}) = 0 \quad (1.3)$$

for  $i = 1, \dots, N-1$  and  $j = 1, \dots, M$ , where we use backward differencing in time and the standard three-point discretization in space:

$$\delta_t U_{ij} := \frac{U_{ij} - U_{i,j-1}}{k_j}, \quad \delta_x^2 U_{ij} := \frac{2}{h_i + h_{i+1}} \left( \frac{U_{i+1,j} - U_{ij}}{h_{i+1}} - \frac{U_{ij} - U_{i-1,j}}{h_i} \right).$$

We also set  $U_{i,0} = \varphi(x_i)$  for  $i = 0, \dots, N$ , and  $U_{0,j} = g_0(t_j)$ ,  $U_{N,j} = g_1(t_j)$  for  $j = 1, \dots, M$ .

Note that the conventional method (1.3), when applied on a uniform mesh in time, might yield incorrect and unstable computed solutions; see Figure 1 (left and centre). Here problem (1.1) was solved with  $f = (2-u)(u-1)u(u+1)$  and  $\varphi = 0.1 + 2x(1-x)$ ,  $g_0 = g_1 = 0.1$ . We observe that  $u(x, 2)$ , which is effectively the steady-state solution, is entirely different from the computed solutions at  $t = 2$ . We also refer the reader to Figure 3 (left), where the numerical method (1.3) is applied to a more complicated problem (5.1) and again yields an incorrect computed solution (which now looks stable and can be easily mistaken for a correct one).

This instability can be explained noting that if  $\varepsilon \ll 1$ , in particular, if  $\varepsilon^2 \ll k_j$ , then the time derivative term  $\varepsilon^2 \delta_t U$ , being  $O(\varepsilon^2/k_j)$ , becomes negligible; thus effectively at each time level we solve a steady-state discrete equation and therefore at each time level we might get any of the multiple steady-state solutions. Furthermore, the space derivative term  $\varepsilon^2 \delta_x^2 U$ , being  $O(\varepsilon^2/(h_i + h_{i+1})^2)$ , might become negligible too, in which case we effectively solve the algebraic equation  $f(x_i, t_j, U_{ij}) = 0$  at each mesh node, where this occurs.




FIG. 1. Computed solutions at  $t = 2$  vs. the exact solution (dashed curve) for various methods;  $\varepsilon = 10^{-2}$ ,  $N = 32$ ,  $M = N^2$ . Left and centre: conventional method (1.3) fails to yield correct computed solutions on the uniform mesh (left), and even if the Shishkin mesh (described in §4.1(b);  $\gamma = 1$ ) is used in space combined with the uniform mesh in time (centre). Right: stabilized method (1.4) with  $\hat{C} = 2$  on the uniform mesh yields a qualitatively correct computed solution.

To stabilize the conventional method (1.3), we generalize it, for some constant  $\hat{C} \geq 0$ , as follows:

$$\hat{\mathcal{J}}^h \hat{U}_{ij} := [\hat{\varepsilon}_j^2 \delta_t - \varepsilon^2 \delta_x^2] \hat{U}_{ij} + f(x_i, t_j, \hat{U}_{ij}) = 0, \quad \hat{\varepsilon}_j^2 = \max\{\varepsilon^2, \hat{C} k_j\}. \quad (1.4)$$

Here, as usual, we set  $\hat{U}_{i,0} = \varphi(x_i)$  for  $i = 0, \dots, N$ , and  $\hat{U}_{0,j} = g_0(t_j)$ ,  $\hat{U}_{N,j} = g_1(t_j)$  for  $j = 1, \dots, M$ . Clearly, (1.3) is a particular case of (1.4) with  $\hat{C} = 0$ . Compared to (1.3), in (1.4) we artificially strengthen the time derivative term, replacing  $\varepsilon^2 \delta_t$  by  $\hat{\varepsilon}_j^2 \delta_t$ , which does not influence the consistency order of the method, but under an appropriate choice of  $\hat{C}$ , always yields a unique computed solution; see Proposition 1.2 below. Furthermore, Figures 1 and 3 illustrate that the instability that we have observed, is indeed cured by switching to the stabilized method (1.4) in which  $\hat{C}$  is chosen using Proposition 1.2.

For uniqueness of solutions of the continuous problem (1.1) and discrete problems (1.3) and (1.4) we have the following results.

**PROPOSITION 1.1 (UNIQUE CONTINUOUS SOLUTION)** Problem (1.1) has at most one solution.

*Proof.* The proof imitates the argument in (Pao, 1992, Theorem 5.1) and we sketch it here for completeness. Suppose (1.1) has two solutions  $u$  and  $\bar{u}$  on  $[0, 1] \times [0, T]$ . Then  $|u|, |\bar{u}| \leq K_1$  and therefore  $f_u \geq -K_2$  in  $[0, 1] \times [0, T] \times [-K_1, K_1]$  for some positive constants  $K_1$  and  $K_2$ , which might depend on  $\varepsilon$  and  $T$ . Using the standard linearization technique and then the transformation  $z := (\bar{u} - u)e^{-tK_2/\varepsilon^2}$ , we get  $\varepsilon^2 [\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2}]z + (K_2 + p)z = 0$  where  $p = p(x, t) = \int_0^1 f_u(x, t, u + s[\bar{u} - u])ds \geq -K_2$ . Since  $z$  vanishes for  $x = 0, 1$  and for  $t = 0$ , by the maximum principle (Protter & Weinberger, 1999, Chapter 3), we have  $z = 0$  for all  $(x, t)$ . Note that this argument relies on  $f_u$  being continuous for all  $u \in \mathbb{R}$  (otherwise, we refer the reader to a solution non-uniqueness example in (Pao, 1992, §1.6)).  $\square$

**PROPOSITION 1.2 (UNIQUE COMPUTED SOLUTION)** Let  $\hat{U}_{ij}$  be a solution of (1.4) and let  $\hat{\varepsilon}_j^2 > C^* k_j$  for some  $C^* \geq 0$ . If  $f_u \geq -C^*$  for all  $x, t, u$ , then  $\hat{U}$  is a unique solution of (1.4). If  $K_1 \leq \hat{U}_{ij} \leq K_2$  for some constants  $K_1$  and  $K_2$ , and  $f_u \geq -C^*$  in  $[0, 1] \times [0, T] \times [K_1, K_2]$ , then  $\hat{U}$  is a unique solution of (1.4) between  $K_1$  and  $K_2$ .

*Proof.* We imitate the proof of (Pao, 1985, Theorem 3.1) (where the case of  $\varepsilon = \hat{\varepsilon}_j = 1$  was considered). We present the proof here, in particular, to reveal the role of  $\hat{\varepsilon}_j$  in the uniqueness condition  $\hat{\varepsilon}_j^2 > C^* k_j$ . Let  $\bar{U}_{ij}$  be another solution of (1.4) and introduce  $Z_{ij} := (\bar{U}_{ij} - \hat{U}_{ij}) \prod_{l=1}^j (1 + k_l \mu_l / \hat{\varepsilon}_l^2)^{-1}$ . A calculation shows that  $\hat{\varepsilon}_j^2 \delta_t Z_{ij} + \mu_j Z_{ij} = (\hat{\varepsilon}_j^2 + k_j \mu_j) [\delta_t (\bar{U}_{ij} - \hat{U}_{ij})] \prod_{l=1}^j (1 + k_l \mu_l / \hat{\varepsilon}_l^2)^{-1}$ . Therefore, applying the standard linearization, we arrive at

$$\frac{\hat{\varepsilon}_j^2}{1 + k_j \mu_j / \hat{\varepsilon}_j^2} \delta_t Z_{ij} - \varepsilon^2 \delta_x^2 Z_{ij} + \left( \frac{\mu_j}{1 + k_j \mu_j / \hat{\varepsilon}_j^2} + p_{ij} \right) Z_{ij} = 0,$$

where  $p_{ij} := \int_0^1 f_u(x_i, t_j, \hat{U}_{ij} + s[\bar{U}_{ij} - \hat{U}_{ij}])ds \geq -C^*$ . As for some constant  $\tilde{C}^* > C^*$  we have  $\hat{\varepsilon}_j^2 \geq \tilde{C}^* k_j$ , then  $\mu_j / (1 + \mu_j k_j \hat{\varepsilon}_j^{-2}) \geq \mu_j / (1 + \mu_j / \tilde{C}^*)$  and choosing  $\mu_j$  sufficiently large, we can always make  $\mu_j / (1 + \mu_j / \tilde{C}^*)$  sufficiently close to  $\tilde{C}^*$ , and therefore, exceeding  $C^*$ . Now  $p_{ij} \geq 0$  and, recalling that  $Z_{ij} = 0$  for  $x_i = 0, 1$  and  $t_j = 0$ , by the discrete maximum principle, we get  $Z_{ij} = 0$  for all  $i, j$ .  $\square$

**REMARK 1.3** To apply Proposition 1.2 to the conventional method (1.3), we have to impose a very restrictive condition  $k_j < \varepsilon^2 / C^*$  on the time step  $k_j$ , which would result in a very inefficient method. In contrast, choosing  $\hat{C}$  sufficiently large in (1.4), we can always ensure a unique computed solution.

The paper is organized as follows. The next §2 presents our assumptions on problem (1.1). In §3 we discuss asymptotic properties of solutions of (1.1) and construct lower and upper solutions. In §4, layer-

adapted meshes for solving (1.1) are described, and discrete analogues of the upper and lower solutions are used to obtain tight upper and lower bounds on the computed solutions. Precise convergence results for the numerical methods (1.3) and (1.4) are then derived on Bakhvalov and Shishkin meshes. In §5, numerical results illustrate the sharpness of our theoretical error estimates. Finally, §6 summarizes our conclusions.

Note that an asymptotic analysis of a version of (1.1) with Neumann boundary conditions, which we partly imitate in §3, was given in (Vasil'eva *et al.*, 1995, §3.2.3). We also refer the reader to asymptotic and numerical analyses for one- and two-dimensional steady-state versions of (1.1) by Fife (1973); Nefedov (1995) and Sun & Stynes (1996); Kopteva & Stynes (2004); Kopteva (2007), respectively.

*Notation.* Throughout this paper we let  $C$  denote a generic positive constant that may take different values in different formulas, but is always independent of  $N$ ,  $M$  and  $\varepsilon$ . A subscripted  $C$  (e.g.,  $C_1$ ) denotes a positive constant that is independent of  $N$ ,  $M$  and  $\varepsilon$  and takes a fixed value. For any two quantities  $w_1$  and  $w_2$ , the notation  $w_1 = O(w_2)$  means  $|w_1| \leq Cw_2$ .

## 2. Assumptions on the continuous problem

We shall examine solutions of (1.1) that exhibit boundary and initial layers. (In general, solutions of (1.1) may also have interior transition layers, which we will consider in a future paper.) As was announced in the introduction, we drop the restrictive global assumption that  $f_u(x, t, u) > 0$  for all  $(x, u) \in [0, 1] \times [0, T] \times \mathbb{R}^1$ , and consider problem (1.1) under the following weaker assumptions.

- It has a *stable reduced solution*, i.e. there exists a sufficiently smooth solution  $u_0$  of (1.2) such that

$$f_u(x, t, u_0(x, t)) > \gamma^2 > 0 \quad \text{for all } (x, t) \in [0, 1] \times [0, T]. \quad (\text{A1})$$

- The boundary conditions satisfy

$$\int_{u_0(l, t)}^v f(l, t, s) ds > 0 \quad \text{for all } v \in (u_0(l, t), g_l(t))', \quad l = 0, 1, \quad t \in [0, T]. \quad (\text{A2})$$

Here the notation  $(a, b)'$  is defined to be  $(a, b)$  when  $a < b$  and  $[b, a)$  when  $a > b$ , while  $(a, b)' = \emptyset$  when  $a = b$ .

- The initial condition is in the domain of attraction of the reduced solution  $u_0$ , i.e. it satisfies

$$s f(x, 0, u_0(x, 0) + s) > 0 \quad \text{for all } s \in (0, \varphi(x) - u_0(x, 0))', \quad x \in [0, 1]. \quad (\text{A3})$$

Note that if  $g_l(t) \approx u_0(l, t)$  for  $l = 0$  or  $l = 1$ , then (A2) follows from (A1) combined with (1.2), while if  $g_l(t) = u_0(l, t)$  at some point  $t \in [0, T]$ , then (A2) does not impose any restriction on  $g_l$  at this point. Similarly, if  $\varphi(x) \approx u_0(x, 0)$ , then (A3) follows from (A1) combined with (1.2), while  $\varphi(x) = u_0(x, 0)$  does not impose any restriction on  $\varphi$  at this point.

Conditions (A1), (A2), (A3) intrinsically arise from the asymptotic analysis of problem (1.1) and guarantee that there exists a unique solution  $u$  of (1.1), which exhibits boundary layers of width  $O(\varepsilon |\ln \varepsilon|)$  at  $x = 0, 1$  and an initial layer of width  $O(\varepsilon^2 |\ln \varepsilon|)$  at  $t = 0$ , while  $u \approx u_0$  in the interior subdomain of  $(0, 1) \times (0, T]$  away from  $x = 0, 1$  and  $t = 0$ ; see Theorem 3.9 for a precise statement. We also refer the reader to Kopteva & Stynes (2004) for a detailed discussion of (A1), (A2) in one dimension, and also to Remark 3.4 on the role of assumption (A3).

We make two further simplifying assumptions to facilitate our presentation. To avoid considering cases, assume that

$$u_0(l, t) \leq g_l(t) \text{ for } l = 0, 1, t \in [0, T]; \quad u_0(x, 0) \leq \varphi(x) \text{ for } x \in [0, 1]. \quad (2.1)$$

To ensure that problem (1.1) has sufficiently smooth solutions, we also impose the *first-order compatibility* conditions  $\varepsilon^2[g'_l(0) - \varphi''(l)] + f(l, 0, \varphi(l)) = 0$  for  $l = 0, 1$ , i.e. at the domain corners  $(0, 0)$  and  $(1, 0)$ . Dropping the  $O(\varepsilon^2)$  terms, we get  $f(l, 0, \varphi(l)) = 0$  for  $l = 0, 1$ . Combining these with (A3), we conclude that

$$\varphi(l) = g_l(0) = u_0(l, 0) \quad \text{for } l = 0, 1. \quad (2.2)$$

Strictly speaking, the terms  $\varepsilon^2[g'_l(0) - \varphi''(l)] = O(\varepsilon^2)$  should remain, and therefore (2.2) should be replaced by a more general relation  $\varphi(l) = g_l(0) = u_0(l, 0) + O(\varepsilon^2)$ . We use (2.2) instead only to simplify our presentation; all our further results apply to this more general case too.

### 3. Asymptotic analysis, upper and lower solutions

We start this section by presenting a standard second-order asymptotic expansion. Furthermore, we shall modify it to construct certain upper and lower solutions that provide tight control on the solutions of our problem (1.1).

We shall use the functions

$$F(x, t, s) := f(x, t, u_0(x, t) + s), \quad \tilde{F}(x, t, s; p) := f(x, t, u_0(x, t) + s) - ps.$$

The perturbed version  $\tilde{F}$  of the function  $F$  is used, with  $|p|$  sufficiently small, in the construction of upper and lower solutions. In the constructions that follow, a tilde will always denote a perturbed function. The perturbed functions always depend on the parameter  $p$ , but we will sometimes not show the explicit dependence. Thus, we will sometimes write  $\tilde{F}(x, t, s)$  for  $\tilde{F}(x, t, s; p)$ . Note that  $\tilde{F}(x, t, 0) = 0$  implies  $\tilde{F}_x(x, t, 0) = 0$ ,  $\tilde{F}_{xx}(x, t, 0) = 0$  and  $\tilde{F}_t(x, t, 0) = 0$ , and therefore we have

$$|\tilde{F}_x(x, t, s)| \leq C|s|, \quad |\tilde{F}_{xx}(x, t, s)| \leq C|s|, \quad |\tilde{F}_t(x, t, s)| \leq C|s|. \quad (3.1)$$

We will occasionally use, for any function  $g$ , the notations

$$g|_a^b = g(b) - g(a), \quad g|_{a;b}^c = g(c) - g(b) - g(a). \quad (3.2)$$

Since  $g(a+b) - g(a) - g(b) + g(0) = abg''(\theta)$ , we see that  $g(0) = 0$  implies  $g|_{a;b}^{a+b} = O(|ab|)$ . Therefore, under this notation,  $\tilde{F}(x, t, 0) = 0$  implies that

$$\tilde{F}(x, t, \cdot)|_{a;b}^{a+b} = O(|ab|). \quad (3.3)$$

Under our assumptions (A1)-(A3), the solution of problem (1.1) exhibits boundary layers near  $x = 0$  and  $x = 1$ , and an initial layer near  $t = 0$ . Since the construction of the layer terms at each of the boundary points is carried out independently of the layer terms at the other boundary point, without loss of generality, we assume throughout this section that

$$u_0(1, t) = g_1(t) \quad \text{for } t \in [0, T], \quad (3.4)$$

which implies that there is no boundary layer at  $x = 1$ . To describe the boundary layer at  $x = 0$  and the initial layer at  $t = 0$ , we shall employ the stretched variables  $\xi := x/\varepsilon$  and  $\tau := t/\varepsilon^2$ .

### 3.1 Solution near the boundary $x = 0$ , boundary-layer functions

In this subsection we construct boundary layer functions associated with the boundary  $x = 0$ ; they use the stretched variable  $\xi = x/\varepsilon$ . Let  $v_0(\xi, t) := \tilde{v}_0(\xi, t; 0)$ , and the functions  $\tilde{v}_0(\xi, t; p)$  and  $v_1(\xi, t)$  be solutions of the equations

$$-\frac{\partial^2 \tilde{v}_0}{\partial \xi^2} + \tilde{F}(0, t, \tilde{v}_0; p) = 0, \quad (3.5a)$$

$$-\frac{\partial^2 v_1}{\partial \xi^2} + v_1 F_s(0, t, v_0) = -\xi F_x(0, t, v_0), \quad (3.5b)$$

where  $\xi > 0$ , subject to the boundary conditions

$$\tilde{v}_0(0, t; p) = g_0(t) - u_0(0, t), \quad v_1(0, t) = 0, \quad \tilde{v}_0(\infty, t; p) = v_1(\infty, t) = 0. \quad (3.5c)$$

Note that the equation for  $\tilde{v}_0$  is a nonlinear autonomous ordinary differential equation, while the equation for  $v_1$  is a linear ordinary differential equation; in these equations,  $t$  and  $p$  appear as parameters. Note also that  $v_1$  is not a perturbed function as it does not depend on  $p$ . Our conditions (A1), (A2) are precisely what is needed to ensure existence and asymptotic properties of  $\tilde{v}_0$  and  $v_1$ . To be more specific, for the solvability and properties of the two problems described by (3.5) we have the following result.

**LEMMA 3.1** Set  $\gamma_L^2 = \min_{t \geq 0} f_u(0, t, u_0(0, t)) > \gamma^2$ , where  $\gamma > 0$  is from (A1). Then there is  $p_0 \in (0, \gamma_L^2)$  such that for all  $|p| \leq p_0$  there exist functions  $\tilde{v}_0(\xi, t; p)$ ,  $v_0(\xi, t)$  and  $v_1(\xi, t)$  which satisfy (3.5). For  $\tilde{v}_0$  and  $v_0$  we have

$$v_0 \geq 0, \quad |v_0 + \varepsilon v_1| \leq Ct, \quad \frac{\partial \tilde{v}_0}{\partial p} \geq 0, \quad \text{for all } \xi, t \geq 0. \quad (3.6)$$

Furthermore, for any arbitrarily small but fixed  $\delta \in (0, \gamma_L - \sqrt{p_0})$ , there is a constant  $C_\delta$  such that

$$\left| \frac{\partial^k \tilde{v}_0}{\partial \xi^k} \right| + \left| \frac{\partial^k v_1}{\partial \xi^k} \right| + \left| \frac{\partial^l \tilde{v}_0}{\partial t^l} \right| + \left| \frac{\partial^l v_1}{\partial t^l} \right| + \left| \frac{\partial \tilde{v}_0}{\partial p} \right| \leq C_\delta e^{-(\gamma_L - \sqrt{p_0} - \delta)\xi} \quad (3.7)$$

for  $\xi, t \geq 0$  and  $k = 0, \dots, 4$ ,  $l = 0, 1, 2$ .

*Proof.* The existence and most of the properties of  $v_0$  and  $\tilde{v}_0$  follow from (Kopteva & Stynes, 2004, Lemma 2.3). For  $v_1$ , we use a result presented in (Fife, 1973, Lemma 2.2) and (Vasil'eva *et al.*, 1995, §2.3.1). In particular, to obtain estimates (3.7), one observes that the derivatives of  $\tilde{v}_0$  and  $v_1$  with respect to  $\xi$  and  $t$ , as well as  $\partial \tilde{v}_0 / \partial p$ , all satisfy linear differential equations with the same differential operator, similar to the one in the equation (3.5b).

We especially elaborate on the proof of  $|v_0 + \varepsilon v_1| \leq Ct$  as its analogues do not appear in the three cited publications. Recall the corner compatibility condition  $g_0(0) - u_0(0, 0) = 0$  from (2.2), which implies  $|g_0(t) - u_0(0, t)| \leq Ct$ . Combining this with  $|v_0(\xi, t)| + |v_1(\xi, t)| \leq C|v_0(0, t)|$  (which follows from the cited analyses of  $v_0$  and  $v_1$ ) and  $v_0(0, t) = g_0(t) - u_0(0, t)$ , yields the desired estimate.  $\square$

For later purposes we shall now obtain two estimates that involve  $\tilde{v}_0$ ,  $v_0$  and  $v_1$ . The first estimate is concerned with the correction  $v_0 + \varepsilon v_1$  to the reduced solution  $u_0$  near  $x = 0$ . We claim that

$$\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (v_0 + \varepsilon v_1) + F(x, t, v_0 + \varepsilon v_1) = O(\varepsilon^2). \quad (3.8)$$

This immediately implies that  $\mathcal{T}(u_0 + v_0 + \varepsilon v_1) = O(\varepsilon^2)$ . Noting that  $(u_0 + v_0 + \varepsilon v_1)|_{x=0} = g_0(t)$  and that  $v_0 + \varepsilon v_1$  is decaying as  $\xi \rightarrow \infty$ , we now expect that  $u_0 + v_0 + \varepsilon v_1$  approximates a solution  $u$  of our problem (1.1) near the boundary  $x = 0$ .

Estimate (3.8) is standard in the asymptotic analysis. It is obtained from  $\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (v_0 + \varepsilon v_1) = -\frac{\partial^2}{\partial \xi^2} (v_0 + \varepsilon v_1) + O(\varepsilon^2)$  combined with (3.5a) and (3.5b), which yield

$$\frac{\partial^2}{\partial \xi^2} (v_0 + \varepsilon v_1) = F(0, t, v_0) + \varepsilon \xi F_x(0, t, v_0) + \varepsilon v_1 F_s(0, t, v_0) = F(\varepsilon \xi, t, v_0 + \varepsilon v_1) + O(\varepsilon^2).$$

Here we also used a Taylor series expansion of  $F(\varepsilon \xi, t, v_0 + \varepsilon v_1)$  in  $\varepsilon$ , in which the quadratic remainder terms were estimated using  $|F_{xx}| \leq C|v_0 + \varepsilon v_1|$  (which follows from (3.1)),  $|F_{ss}| + |F_{xs}| \leq C$ , and then  $(\xi^2 + 1)(|v_0| + |v_1|) \leq C$  (which follows from (3.7)). Thus (3.8) is established.

Our second auxiliary estimate is for  $\tilde{v}_0 - v_0$ :

$$\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (\tilde{v}_0 - v_0) = -F(x, t, \cdot) \Big|_{v_0 + \varepsilon v_1}^{\tilde{v}_0 + \varepsilon v_1} + p v_0 + O(\varepsilon^2 + p^2). \quad (3.9)$$

It follows from  $\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (\tilde{v}_0 - v_0) = -\frac{\partial^2}{\partial \xi^2} (\tilde{v}_0 - v_0) + O(\varepsilon^2)$  combined with (3.5a), which implies

$$\frac{\partial^2}{\partial \xi^2} (\tilde{v}_0 - v_0) = F(0, t, \cdot) \Big|_{v_0}^{\tilde{v}_0} + p \tilde{v}_0 = F(x, t, \cdot) \Big|_{v_0 + \varepsilon v_1}^{\tilde{v}_0 + \varepsilon v_1} - x F_x(\hat{x}, t, \cdot) \Big|_{v_0 + \varepsilon \delta}^{\tilde{v}_0 + \varepsilon \delta} - \varepsilon v_1 F_s(\hat{x}, t, \cdot) \Big|_{v_0 + \varepsilon \delta}^{\tilde{v}_0 + \varepsilon \delta} + p \tilde{v}_0.$$

Recalling that  $x = \varepsilon \xi$  and noting that, by the estimate for  $\frac{\partial \tilde{v}_0}{\partial p}$  in (3.7), we have  $(1 + \xi)|\tilde{v}_0 - v_0| = O(p)$ , yields (3.9).

### 3.2 Solution near $t = 0$ , initial-layer functions

In this subsection we construct initial-layer functions to describe the solution near  $t = 0$ ; they use the stretched variable  $\tau = t/\varepsilon^2$ . Let  $w_0(x, \tau) := \tilde{w}_0(x, \tau; 0)$ , and the function  $\tilde{w}_0(x, \tau; p)$  be a solution of the initial-value problem

$$\frac{\partial \tilde{w}_0}{\partial \tau} = -\tilde{F}(x, 0, \tilde{w}_0; p) \quad \text{for } \tau > 0, \quad \tilde{w}_0(x, 0; p) = \varphi(x) - u_0(x, 0). \quad (3.10a)$$

Since  $w_0$  and  $\tilde{w}_0$  describe a correction to  $u_0(x, t)$  for small values of  $t$ , we look for a solution of (3.10a) that satisfies an additional condition

$$\tilde{w}_0(x, \infty; p) = 0. \quad (3.10b)$$

Here  $x \in [0, 1]$  and  $p$  appear as parameters.

For each fixed  $x$  and  $p$ , problem (3.10) is a particular case of the auxiliary initial value problem

$$\frac{d}{d\tau} \omega = -\phi(\omega) \quad \text{for } \tau > 0, \quad \omega(0) = \omega_0 \geq 0, \quad \omega(\infty) = 0, \quad (3.11)$$

for which we have the following result

LEMMA 3.2 Let a sufficiently smooth function  $\phi$  satisfy

$$\phi(0) = 0, \quad \phi'(0) > 0, \quad \phi(s) > 0 \quad \text{for all } s \in (0, \omega_0]. \quad (3.12)$$

(i) Then problem (3.11) has a solution  $0 \leq \omega \leq \omega_0$ , and for any arbitrarily small but fixed  $\delta \in (0, \phi'(0))$ , there is a constant  $C_\delta$  such that

$$|\omega| + |\omega'| + |\omega''| \leq \omega_0 C_\delta e^{-[\phi'(0) - \delta]\tau} \quad \text{for } \tau \geq 0. \quad (3.13)$$

(ii) Set  $\hat{\omega} := \omega/\omega_0$  if  $\omega_0 > 0$ , or  $\hat{\omega} := e^{-\phi'(0)\tau}$  if  $\omega_0 = 0$ . Then the related linear problem

$$\frac{d}{d\tau}\chi + \chi\phi'(\omega) = \psi(\tau) \quad \text{for } \tau > 0, \quad \chi(0) = \chi_0, \quad \chi(\infty) = 0, \quad (3.14)$$

where  $|\psi(\tau)| \leq C(1 + \tau^m)\hat{\omega}(\tau)$  for some  $m \geq 0$ , has a solution such that  $|\chi(\tau)| \leq C(\chi_0 + 1 + \tau^{m+1})\hat{\omega}(\tau)$ . If we also have  $\chi_0 = 0$  and  $\psi \geq 0$ , then  $\chi \geq 0$  for all  $\tau \geq 0$ .

*Proof.* (i) If  $\omega_0 = 0$ , then  $\omega(\tau) = 0$  for all  $\tau$  and the assertion follows. Otherwise, if  $\omega_0 > 0$ , consider the phase plane  $(\omega, \omega')$  for the equation  $\omega' = -\phi(\omega)$ . By (3.12), there is a trajectory that leaves the point  $(\omega_0, -\phi(\omega_0))$  and enters the point  $(0, 0)$ , which is a fixed point for this equation. Furthermore, since  $\phi(\omega) > 0$  for all  $\omega \in (0, \omega_0]$ , this entire trajectory will lie in the quarter plane  $\{\omega > 0, \omega' < 0\}$ ; therefore the corresponding solution  $\omega(\tau)$  is positive and decreasing to 0. It remains to show that the solution trajectory enters  $(0, 0)$  as  $\tau \rightarrow \infty$ , and also the exponential decay estimates (3.13). Note that for any  $\delta \in (0, \phi'(0))$ , there exists  $s_\delta \in (0, \omega_0)$  such that  $[\phi'(0) - \delta]s \leq \phi(s) \leq [\phi'(0) + \delta]s$  for all  $s \in [0, s_\delta]$ . Furthermore, there exists  $\tau_\delta > 0$  such that  $\omega(\tau_\delta) = s_\delta$  (otherwise, if  $\omega(\tau) > s_\delta$  for all  $\tau$ , then  $\omega'(\tau) \leq -C$  for some positive constant  $C$ , which yields a contradiction  $\omega(\infty) = -\infty$  with (3.11)). Thus for all  $\tau \geq \tau_\delta$  we have  $[\phi'(0) - \delta]\omega \leq \omega' \leq [\phi'(0) + \delta]\omega$ , which implies that  $e^{-[\phi'(0) + \delta]\tau} \leq \omega(\tau)/\omega(\tau_\delta) \leq e^{-[\phi'(0) - \delta]\tau}$  for  $\tau \geq \tau_\delta$ . The estimates for  $\omega$  and  $\omega'$  in (3.13) follow immediately as  $\omega(\tau_\delta) < \omega_0$ . Finally, the estimate for  $\omega''$  in (3.13) is obtained noting that  $\omega'' = -\omega'\phi'(\omega)$ .

(ii) To solve (3.14), note that the corresponding homogeneous equation  $\frac{d}{d\tau}\theta + \theta\phi'(\omega) = 0$  has a positive solution  $\theta$  such that  $\theta(0) = 1$ . If  $\omega_0 > 0$ , then we recall from part (i) that  $\omega' < 0$  and  $C^{-1} \leq |\omega'|/\omega \leq C$  and thus choose  $\theta := \omega'/\omega'(0) > 0$  so that  $C^{-1} \leq \theta/\hat{\omega} \leq C$ ; otherwise, if  $\omega_0 = 0$  and thus  $\omega = 0$ , then, by (3.12), we have  $\phi'(\omega) = \phi'(0) > 0$  and so choose  $\theta(\tau) := e^{-\phi'(0)\tau} = \hat{\omega}$ . Now, the unique solution of (3.14) is given by

$$\chi(\tau) = \chi_0 \theta(\tau) + \theta(\tau) \int_0^\tau \frac{\psi(\tau')}{\theta(\tau')} d\tau',$$

where  $|\psi(\tau)| \leq C(1 + \tau^m)\theta(\tau)$ . The desired assertions follow.  $\square$

We now apply Lemma 3.2 to problem (3.10) as follows.

LEMMA 3.3 Set  $\gamma_0^2 := \min_{x \in [0,1]} f_u(x, 0, u_0(x, 0)) > \gamma^2$ , where  $\gamma > 0$  is from (A1). Then there is  $p_0 \in (0, \gamma_0^2)$  such that for all  $|p| \leq p_0$ , problem (3.10) has a solution  $\tilde{w}_0(x, \tau; p)$ . For  $\tilde{w}_0$  and  $w_0$  we have

$$0 \leq w_0 \leq Cx, \quad \frac{\partial \tilde{w}_0}{\partial p} \geq 0, \quad \text{for all } x \in [0, 1], \tau \geq 0. \quad (3.15)$$

Furthermore, for any arbitrarily small but fixed  $\delta \in (0, \gamma_0^2 - p_0)$ , there is a constant  $C_\delta$  such that

$$\left| \frac{\partial^l \tilde{w}_0}{\partial \tau^l} \right| + \left| \frac{\partial^k \tilde{w}_0}{\partial x^k} \right| + \left| \frac{\partial \tilde{w}_0}{\partial p} \right| \leq C_\delta e^{-(\gamma_0^2 - p_0 - \delta)\tau}. \quad (3.16)$$

for  $x \in [0, 1]$ ,  $\tau \geq 0$  and  $k = 0, \dots, 4$ ,  $l = 0, 1, 2$ .

*Proof.* For each fixed  $x$  and  $p$ , problem (3.10) is a particular case of the auxiliary problem (3.11) with a solution  $\omega := \tilde{w}_0$ , the initial condition  $\omega_0 := \varphi(x) - u_0(x, 0)$ , and the right-hand side function  $\phi(s) := \tilde{F}(x, 0, s; p) = f(x, 0, u_0(x, 0) + s) - ps$ , for which we have  $\phi'(s) = f_u(x, 0, u_0(x, 0) + s) - p$ . Note that (A1) implies that  $\phi(0) = 0$  and  $\phi'(0) \geq \gamma_0^2 - |p| > 0$ , while (A3) combined with (2.1) yields  $\phi(s) > 0$  for all  $s \in (0, \omega_0]$  provided that  $p_0$  is chosen sufficiently small. Thus the hypotheses (3.12) of



Lemma 3.2 are satisfied. Now Lemma 3.2(i) implies existence of a solution  $0 \leq \tilde{w}_0 \leq \varphi(x) - u_0(x, 0)$  and the estimates for  $\frac{\partial^l}{\partial \tau^l} \tilde{w}_0$ , where  $l = 0, 1, 2$ , in (3.16). Next, the bound  $0 \leq w_0 \leq Cx$  in (3.15) is obtained from  $w_0 = \tilde{w}_0|_{p=0}$  and  $0 \leq \tilde{w}_0 \leq \varphi(x) - u_0(x, 0)$ , noting that the corner compatibility condition  $\varphi(0) - u_0(0, 0) = 0$  from (2.2) implies  $\varphi(x) - u_0(x, 0) \leq Cx$ .

It remains now to estimate the functions  $\frac{\partial}{\partial p} \tilde{w}_0$  and  $\frac{\partial^k}{\partial x^k} \tilde{w}_0$  for  $k = 1, \dots, 4$ . Differentiating the equation in (3.10a) with respect to  $p$ , or  $k$  times with respect to  $x$ , we see that these functions are solutions of the initial value problem (3.14) with  $\phi'(\omega) = \phi'(\tilde{w}_0) = \tilde{F}_s(x, 0, \tilde{w}_0)$  and various right-hand sides and initial data. In particular, we have  $\psi := \tilde{w}_0$  and  $\chi_0 := 0$  for  $\chi = \frac{\partial}{\partial p} \tilde{w}_0$ , which, by Lemma 3.2(ii), implies that  $\frac{\partial}{\partial p} \tilde{w}_0 \geq 0$  and the estimate for this function in (3.16). Furthermore, for  $\chi = \frac{\partial}{\partial x} \tilde{w}_0$  we use  $\psi := -\tilde{F}_x(x, 0, \tilde{w}_0)$  (for which we have  $|\psi| \leq C\tilde{w}_0$ , by (3.1)) and  $\chi_0 := \varphi'(x) - u_{0,x}(x, 0)$ ; now the estimate for  $\frac{\partial}{\partial x} \tilde{w}_0$  in (3.16) is obtained by again applying Lemma 3.2(ii). The remaining bounds in (3.16) are obtained similarly, by evaluating the functions  $\psi$  and  $\chi_0$  corresponding to  $\chi = \frac{\partial^k}{\partial x^k} \tilde{w}_0$  with  $k > 1$ , and then applying Lemma 3.2(ii).  $\square$

For later purposes we shall now obtain two estimates that involve  $\tilde{w}_0$  and  $w_0$ . The first estimate is concerned with the correction  $w_0$  to the reduced solution  $u_0$  near  $t = 0$ . We claim that

$$\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] w_0 + F(x, t, w_0) = O(\varepsilon^2). \quad (3.17)$$

This immediately implies that  $\mathcal{T}(u_0 + w_0) = O(\varepsilon^2)$ . Noting that  $(u_0 + w_0)|_{t=0} = \varphi(x)$  and that  $w_0$  is decaying as  $\tau \rightarrow \infty$ , we expect that  $u_0 + w_0$  approximates a solution  $u$  of our problem (1.1) near  $t = 0$ .

Estimate (3.17) is standard in asymptotic analysis. It is obtained noting that  $\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] w_0 = \frac{\partial}{\partial \tau} w_0 + O(\varepsilon^2)$  and then recalling (3.10a), which yields  $\frac{\partial}{\partial \tau} w_0 = -F(x, 0, w_0) = -F(x, t, w_0) + O(\varepsilon^2)$ . Here we also used a Taylor series expansion of  $F(x, t, w_0)$  in  $t$ , in which the linear remainder term  $tF_t(x, \hat{t}, w_0)$  was estimated combining  $t = \varepsilon^2 \tau$  with  $|F_t| \leq Cw_0$  (which follows from (3.1)) and then invoking (3.16). Thus (3.17) is established.

Our second auxiliary estimate is for  $\tilde{w}_0 - w_0$ :

$$\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (\tilde{w}_0 - w_0) = -F(x, t, \cdot) \Big|_{w_0}^{\tilde{w}_0} + pw_0 + O(\varepsilon^2 + p^2). \quad (3.18)$$

It follows from  $\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (\tilde{w}_0 - w_0) = \frac{\partial}{\partial \tau} (\tilde{w}_0 - w_0) + O(\varepsilon^2)$  combined with (3.10a), which implies

$$\frac{\partial}{\partial \tau} (\tilde{w}_0 - w_0) = -F(x, 0, \cdot) \Big|_{w_0}^{\tilde{w}_0} + p\tilde{w}_0 = -F(x, t, \cdot) \Big|_{w_0}^{\tilde{w}_0} + tF_t(x, \hat{t}, \cdot) \Big|_{w_0}^{\tilde{w}_0} + p\tilde{w}_0.$$

Recalling that  $t = \varepsilon^2 \tau$  and noting that, by the bound for  $\frac{\partial \tilde{w}_0}{\partial p}$  in (3.16), we have  $(1 + \tau)|\tilde{w}_0 - w_0| = O(p)$ , yields (3.18).

**REMARK 3.4** Assumption (A3) is necessary for existence of the initial-layer functions  $w_0$  and  $\tilde{w}_0$ , which are solutions of problem (3.10). This follows from (3.10) being a particular case of problem (3.11), as an extension of the phase plane analysis used in the proof of Lemma 3.2, shows that, if  $|\phi'(s)| \leq C$  for all  $s \in [0, \omega_0]$ , then our conditions (3.12), with  $\phi'(0) > 0$  relaxed to  $\phi'(0) \geq 0$ , are necessary for problem (3.11) having a solution.

### 3.3 First-order asymptotic expansion

In the previous subsections we have defined the boundary-layer functions  $v_0$  and  $v_1$  and the initial-layer function  $w_0$ . In this subsection, these functions and the reduced solution  $u_0$  are assembled in the following first-order asymptotic expansion for our problem (1.1):

$$u_{\text{as}}(x, t) := u_0(x, t) + [v_0(\xi, t) + \varepsilon v_1(\xi, t)] + w_0(x, \tau). \quad (3.19)$$

Note that no corner functions are needed in the above asymptotic expansion due to the compatibility conditions (2.2). Indeed, examining problems (3.5) for  $v_0$  and  $v_1$ , in view of (2.2) with  $l = 0$ , yields  $v_0(\xi, 0) = v_1(\xi, 0) = 0$  for all  $\xi \geq 0$ ; similarly, examining problem (3.10) for  $w_0$  in view of (2.2), yields  $w_0(0, \tau) = w_0(1, \tau) = 0$  for all  $\tau \geq 0$ . Therefore, we get

$$u_{\text{as}}(x, 0) = \varphi(x), \quad u_{\text{as}}(0, t) = g_0(t), \quad u_{\text{as}}(1, t) = g_1(t) + O(\varepsilon^2), \quad (3.20)$$

or in other words,  $u_{\text{as}}(x, 0) = u(x, 0)$  and  $|u_{\text{as}}(l, t) - u(l, t)| = O(\varepsilon^2)$  at the boundary points  $l = 0, 1$ . It should be noted that the last relation in (3.20) follows from  $u_{\text{as}}(1, t) = u_0(1, t) + (v_0 + \varepsilon v_1)|_{\xi=1/\varepsilon}$  combined with our assumption (3.4) and the estimate  $|v_0 + \varepsilon v_1| \leq C_\delta e^{-(\nu-\delta)/\varepsilon} \leq C\varepsilon^2$  for  $\xi = 1/\varepsilon$ , for which we invoked (3.7).

Furthermore, we have the following standard result for  $\mathcal{T}u_{\text{as}}$ .

LEMMA 3.5 The asymptotic expansion  $u_{\text{as}}$  from (3.19) satisfies  $\mathcal{T}u_{\text{as}} = O(\varepsilon^2)$ .

*Proof.* First we combine  $\varepsilon^2 [\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2}] u_0 = O(\varepsilon^2)$  with (3.8) and (3.17) and, using notation (3.2), get

$$\mathcal{T}u_{\text{as}} = \varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] u_{\text{as}} + F(x, t, v_0 + \varepsilon v_1 + w_0) = F(x, t, \cdot) \Big|_{v_0 + \varepsilon v_1; w_0}^{(v_0 + \varepsilon v_1) + w_0} + O(\varepsilon^2).$$

By (3.3), this yields

$$|\mathcal{T}u_{\text{as}}| \leq C |(v_0 + \varepsilon v_1)w_0| + O(\varepsilon^2) \leq C t e^{-(\nu_0 - \delta)\tau} + O(\varepsilon^2) \leq C\varepsilon^2.$$

Here we estimated  $|v_0 + \varepsilon v_1|$  using (3.6) and  $w_0$  using (3.16), and also invoked  $t = \varepsilon^2 \tau$ .  $\square$

### 3.4 Modified asymptotic expansion, existence of a solution between upper and lower solutions

In this section we construct upper and lower solutions, and therefore, prove an existence of a solution in an  $O(\varepsilon^2)$  neighbourhood of our asymptotic expansion. The upper and lower solutions are obtained by perturbing our asymptotic expansion (3.19), in which we replace the boundary- and initial-layer functions  $v_0$  and  $w_0$  by their perturbed versions  $\tilde{v}_0$  and  $\tilde{w}_0$ , and then add the term  $C_0 p$ , as follows:

$$\beta(x, t; p) := u_0(x, t) + [\tilde{v}_0(\xi, t; p) + \varepsilon v_1(\xi, t)] + \tilde{w}_0(x, \tau; p) + C_0 p. \quad (3.21)$$

Occasionally we shall use an alternative equivalent representation

$$\beta(x, t; p) = u_{\text{as}} + V + W + C_0 p, \quad \text{where } V := \tilde{v}_0 - v_0, \quad W := \tilde{w}_0 - w_0. \quad (3.22)$$

Note that for  $V$  and  $W$  here, by the estimates for  $\frac{\partial}{\partial p} \tilde{v}_0$  and  $\frac{\partial}{\partial p} \tilde{w}_0$  in (3.7) and (3.16), we get

$$(1 + \xi)|V| \leq Cp, \quad (1 + \tau)|W| \leq Cp. \quad (3.23)$$

LEMMA 3.6 For the function  $\beta(x, t; p)$  of (3.21) we have  $\beta = u_{\text{as}} + O(p)$ . Furthermore, if  $p \geq 0$ , then

$$\beta(x, t; -p) \leq u_{\text{as}} - C_0 p \leq u_{\text{as}} + C_0 p \leq \beta(x, t; p) \quad \text{for all } (x, t) \in [0, 1] \times [0, T]. \quad (3.24)$$

*Proof.* The first assertion immediately follows from (3.22) and (3.23). Noting that  $u_{\text{as}}(x, t) = \beta(x, t; 0)$  and then recalling the bounds  $\partial \bar{v}_0 / \partial p \geq 0$  and  $\partial \bar{w}_0 / \partial p \geq 0$  from (3.6) and (3.15), yields the second assertion (3.24).  $\square$

Furthermore, for  $\mathcal{T}\beta$  we get the following result.

LEMMA 3.7 For all  $(x, t) \in (0, 1) \times (0, T]$  we have

$$\mathcal{T}\beta = C_0 p f_u(x, t, u_0) + p[1 + C_0 \lambda](v_0 + w_0) + O(\varepsilon^2 + p^2),$$

where  $\lambda = \lambda(x, t) := f_{uu}(x, t, u_0 + \vartheta[v_0 + w_0])$  for some  $\vartheta = \vartheta(x, t) \in (0, 1)$ .

*Proof.* By Lemma 3.5, we have  $\mathcal{T}u_{\text{as}} = O(\varepsilon^2)$ . Thus it suffices to investigate  $\mathcal{T}\beta - \mathcal{T}u_{\text{as}}$ , for which, by (3.22), we have

$$\mathcal{T}\beta - \mathcal{T}u_{\text{as}} = \varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (V + W) + f(x, t, \cdot) \Big|_{u_{\text{as}}}^{\beta}. \quad (3.25)$$

Now, recalling (3.9) and (3.18) yields

$$\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (V + W) = -F(x, t, \cdot) \Big|_{v_0 + \varepsilon v_1}^{\bar{v}_0 + \varepsilon v_1} - F(x, t, \cdot) \Big|_{w_0}^{\bar{w}_0} + p[v_0 + w_0] + O(\varepsilon^2 + p^2),$$

and therefore, using Taylor series expansions combined with  $V^2 + W^2 \leq Cp^2$  (see (3.23)), we get

$$\varepsilon^2 \left[ \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right] (V + W) = -VF_s(x, t, v_0 + \varepsilon v_1) - WF_s(x, t, w_0) + p(v_0 + w_0) + O(\varepsilon^2 + p^2). \quad (3.26)$$

Similarly, we obtain

$$\begin{aligned} f(x, t, \cdot) \Big|_{u_{\text{as}}}^{u_{\text{as}} + V + W} &= F(x, t, \cdot) \Big|_{v_0 + \varepsilon v_1 + w_0}^{v_0 + \varepsilon v_1 + w_0 + V + W} = [V + W]F_s(x, t, v_0 + \varepsilon v_1 + w_0) + O(p^2) \\ &= VF_s(x, t, v_0 + \varepsilon v_1) + WF_s(x, t, w_0) + O(|Vw_0| + |W(v_0 + \varepsilon v_1)| + p^2). \end{aligned} \quad (3.27)$$

Next, note that invoking  $\varepsilon v_1 + (V + W) = O(\varepsilon + p)$ , we get

$$\begin{aligned} f(x, t, \cdot) \Big|_{u_{\text{as}} + V + W}^{\beta} &= f(x, t, \cdot) \Big|_{u_{\text{as}} + V + W}^{u_{\text{as}} + V + W + C_0 p} = C_0 p [f_u(x, t, u_0 + v_0 + w_0) + O(\varepsilon + p)] \\ &= C_0 p [f_u(x, t, u_0) + \lambda(v_0 + w_0)] + O(\varepsilon^2 + p^2), \end{aligned} \quad (3.28)$$

where  $\lambda = \lambda(x, t) := f_{uu}(x, t, u_0 + \vartheta[v_0 + w_0])$  for some  $\vartheta = \vartheta(x, t) \in (0, 1)$ .

Combining relations (3.25), (3.26), (3.27), (3.28) with  $\mathcal{T}u_{\text{as}} = O(\varepsilon^2)$ , we arrive at

$$\mathcal{T}\beta = C_0 p f_u(x, t, u_0) + p[1 + C_0 \lambda](v_0 + w_0) + O(|Vw_0| + |W(v_0 + \varepsilon v_1)|) + O(\varepsilon^2 + p^2).$$

The desired assertion follows by invoking  $|Vw_0| + |W(v_0 + \varepsilon v_1)| \leq C(x|V| + t|W|) = O(\varepsilon p)$ . Here we estimated  $|w_0|$  and  $|v_0 + \varepsilon v_1|$  using (3.15) and (3.6), and then recalled  $x = \varepsilon \xi$  and  $\tau = \varepsilon^2 \tau$  combined with  $\xi|V| \leq Cp$  and  $\tau|W| \leq Cp$  from (3.23).  $\square$

COROLLARY 3.8 There are  $C_0 > 0$  and  $C_1 > 0$  such that for all  $|p| \leq p_0$  we have

$$\begin{aligned} \mathcal{T}\beta &\geq C_0 p \gamma^2 - C_1(\varepsilon^2 + p^2), & \text{if } p > 0, \\ \mathcal{T}\beta &\leq -C_0 |p| \gamma^2 + C_1(\varepsilon^2 + p^2), & \text{if } p < 0. \end{aligned}$$

*Proof.* Recall (A1) and the estimates  $v_0 \geq 0$ ,  $w_0 \geq 0$  from (3.6), (3.15). Now choose  $0 < C_0 \leq |\lambda(x, t)|^{-1}$  for all  $x$  and  $t$  so that  $1 + C_0 \lambda \geq 0$ .  $\square$

Now we are ready to establish existence of a unique solution of (1.1) that lies in an  $O(\varepsilon^2)$  neighbourhood of our asymptotic expansion.

THEOREM 3.9 There is a sufficiently small  $\varepsilon_0 > 0$  such that for all  $\varepsilon \leq \varepsilon_0$ , there exists a unique solution  $u$  of problem (1.1). Furthermore, for this solution we have  $|u(x, t) - u_{\text{as}}(x, t)| \leq C\varepsilon^2$  for all  $(x, t) \in [0, 1] \times [0, T]$ .

*Proof.* Set  $\bar{p} = C_2 \varepsilon^2$ , where  $C_2 \geq 2C_1/(C_0 \gamma^2)$  so that  $C_0 \bar{p} \gamma^2 \geq 2C_1 \varepsilon^2$ . Then, by Corollary 3.8, for  $\varepsilon \leq 1/C_2$  we get  $\bar{p} \leq \varepsilon$  so  $C_1(\varepsilon^2 + p^2) \leq 2C_1 \varepsilon^2$  and therefore

$$\mathcal{T}\beta(x, t; -\bar{p}) \leq 0 \leq \mathcal{T}\beta(x, t; \bar{p}). \quad (3.29a)$$

Furthermore, in view of (3.24), choosing  $C_2$  sufficiently large so that  $C_0 \bar{p} = C_0 C_2 \varepsilon^2$  dominates the term  $O(\varepsilon^2)$  in (3.20), yields

$$\beta(x, 0; -\bar{p}) \leq \varphi(x) \leq \beta(x, 0; \bar{p}), \quad \beta(l, t; -\bar{p}) \leq g_l(t) \leq \beta(l, t; \bar{p}) \quad \text{for } l = 0, 1. \quad (3.29b)$$

By (3.24), we also have

$$\beta(x, t; -\bar{p}) \leq \beta(x, t; \bar{p}). \quad (3.29c)$$

Comparing (3.29) with (1.1), we see that  $\beta(x, t; -\bar{p})$  and  $\beta(x, t; \bar{p})$  are ordered lower and upper solutions, respectively, for problem (1.1) (sometimes they are called ordered sub- and super-solutions); see Pao (1992). Now, applying (Pao, 1992, Theorem 5.1) yields existence of a solution  $u$  between  $\beta(x, t; -\bar{p})$  and  $\beta(x, t; \bar{p})$ :

$$\beta(x, t; -\bar{p}) \leq u(x, t) \leq \beta(x, t; \bar{p}).$$

Furthermore, Proposition 1.1 implies that this is a unique solution. Since, by Lemma 3.6, we have  $\beta(x, t; \pm \bar{p}) = u_{\text{as}} + O(\bar{p}) = u_{\text{as}} + O(\varepsilon^2)$ , then  $|u - u_{\text{as}}| \leq C\varepsilon^2$ .  $\square$

#### 4. Analysis of the numerical method

In this section we investigate the numerical method (1.4); note that our results also apply to a more conventional numerical method (1.3) as it is a particular case of (1.4) with  $\hat{C} = 0$ .

We make a further simplifying assumption to facilitate our presentation. Throughout this section we take

$$\varepsilon \leq C(N^{-1} + M^{-1/2}). \quad (4.1)$$

This is not a practical restriction, and from a theoretical viewpoint the analysis of a nonlinear problem such as (1.1) would be very different if  $\varepsilon$  were not small. Furthermore, by invoking higher-order asymptotic expansions (compared to (3.19)), condition (4.1) can be relaxed to  $\varepsilon \leq C(N^{-\delta} + M^{-\delta/2})$  for any arbitrarily small but fixed  $\delta \in (0, 1]$ .

#### 4.1 Layer-adapted meshes, truncation error

We shall consider discrete problems (1.3) and (1.4) on two popular layer-adapted meshes, which have been shown to yield convergence of various numerical methods uniformly with respect to the singular perturbation parameter(s). The meshes are presented for the general case when the solution of problem (1.1) has boundary layers both at  $x = 0$  and  $x = 1$  and also an initial layer at  $t = 0$ ; see Figure 2. For convenience, we nevertheless continue our analysis in this section under assumption (3.4).

4.1 (a) *Bakhvalov mesh* first appeared in Bakhvalov (1969); we also refer the reader to Roos *et al.* (2008). The mesh points  $(x_i, t_j)$  are defined as  $x_i = x(i/N)$  and  $t_j = t(j/M)$ , where the mesh-generating functions  $x(\cdot), t(\cdot) \in C[0, 1]$  are given by

$$x(\xi) = \begin{cases} \frac{2\varepsilon}{\gamma} \ln \frac{1}{1-4\xi} & \text{for } \xi \in [0, \theta], \\ \frac{1}{2} - d(\frac{1}{2} - \xi) & \text{for } \xi \in (\theta, \frac{1}{2}], \\ 1 - x(1 - \xi) & \text{for } \xi \in (\frac{1}{2}, 1], \end{cases} \quad t(\eta) = \begin{cases} \frac{\varepsilon^2}{\gamma^2} \ln \frac{1}{1-2\eta} & \text{for } \eta \in [0, \theta_0], \\ T - d_0(1 - \eta) & \text{for } \eta \in (\theta_0, 1]. \end{cases}$$

Here  $\theta = 1/4 - C_3\varepsilon$  and  $\theta_0 = 1/2 - C_4\varepsilon^2$  for some positive constants  $C_3$  and  $C_4$ ; and  $d$  and  $d_0$  are chosen so that  $x(\xi)$  and  $t(\eta)$  are continuous at  $\xi = \theta$  and  $\eta = \theta_0$  respectively. These definitions of  $x(\xi)$  and  $t(\eta)$  are valid only for  $\varepsilon \leq \frac{1}{8} \min\{\gamma, 2C_3^{-1}\}$  and  $\varepsilon^2 \leq \frac{1}{2} \min\{\gamma^2 T, C_4^{-1}\}$ , respectively, which is not a practical restriction (otherwise, we set  $x(\xi) = \xi$  and/or  $t(\eta) = T\eta$  and get a uniform mesh in the  $x$ - and/or  $t$ -direction). Note also that for a certain choice of  $C_3$  and  $C_4$ , one obtains the original Bakhvalov mesh, for which  $x(\cdot), t(\cdot) \in C^1[0, 1]$ .

4.1 (b) *Shishkin mesh*; see Shishkin (1992); Miller *et al.* (1996). This mesh is constructed as follows. Let  $N/4$  and  $M/2$  be positive integers and set

$$\sigma := \min\left\{\frac{2\varepsilon}{\gamma} \ln N, \frac{1}{4}\right\}, \quad \sigma_0 := \min\left\{\frac{\varepsilon^2}{\gamma^2} \ln M, \frac{T}{2}\right\}. \quad (4.2)$$

Now the piecewise uniform mesh  $\{x_i\}_{i=0}^N$  is obtained by dividing the intervals  $[0, \sigma]$ ,  $[\sigma, 1 - \sigma]$  and  $[1 - \sigma, 1]$  into  $N/4$ ,  $N/2$  and  $N/4$  equidistant subintervals, respectively. Similarly the piecewise uniform mesh  $\{t_j\}_{j=0}^M$  is obtained by dividing each of the intervals  $[0, \sigma_0]$  and  $[\sigma_0, T]$  into  $M/2$  equidistant subintervals. In practice, one usually has  $\sigma \ll 1$  and  $\sigma_0 \ll 1$ , so the  $x$ -mesh is coarse on  $[\sigma, 1 - \sigma]$  and fine otherwise, while the  $t$ -mesh is coarse on  $[\sigma_0, T]$  and fine on  $[0, \sigma_0]$ .

For the truncation error  $\hat{\mathcal{J}}^h \beta - \mathcal{J} \beta$  of  $\hat{\mathcal{J}}^h$  from (1.4) on these meshes we have the following estimate.

LEMMA 4.1 Let  $\beta(x, t) = \beta(x, t; p)$  be defined by (3.21), and let the mesh  $\{(x_i, t_j)\}$  be either the Bakhvalov mesh of §4.1(a), or the Shishkin mesh of §4.1(b). Then for all  $|p| \leq p_0$ , where  $p_0$  is a sufficiently small constant, we have

$$|\hat{\mathcal{J}}^h \beta(x_i, t_j; p) - \mathcal{J} \beta(x_i, t_j; p)| \leq C(N^{-2} \ln^{2m} N + M^{-1} \ln^m M),$$

where  $m = 0$  for the Bakhvalov mesh (a) and  $m = 1$  for the Shishkin mesh (b).

*Proof.* Choose  $p_0$  in Lemmas 3.1 and 3.3 sufficiently small so that  $\gamma_L - \sqrt{p_0} > \gamma$  and  $\gamma_0^2 - p_0 > \gamma^2$ ; next, choose  $\delta$  in (3.7) and (3.16) sufficiently small so that  $\gamma_L - \sqrt{p_0} - \delta \geq \gamma$  and  $\gamma_0^2 - p_0 - \delta \geq \gamma^2$ .

Next, note that  $\hat{\mathcal{J}}^h \beta - \mathcal{J} \beta = [\hat{\varepsilon}^2 \delta_t - \varepsilon^2 \frac{\partial}{\partial t}] \tilde{w}_0 - \varepsilon^2 [\delta_x^2 - \frac{\partial^2}{\partial x^2}] (\tilde{v}_0 + \varepsilon v_0) + O(\varepsilon^2)$ . By (3.7), imitating the arguments in (Kopteva & Stynes, 2004, §3.4), we get  $\varepsilon^2 [\delta_x^2 - \frac{\partial^2}{\partial x^2}] (\tilde{v}_0 + \varepsilon v_0) = O(N^{-2} \ln^{2m} N)$ . Recalling (4.1), we observe that to establish the desired estimate it now remains to show that

$$R_1 := \varepsilon^2 [\delta_t - \frac{\partial}{\partial t}] \tilde{w}_0 = O(M^{-1} \ln^m M), \quad R_2 := (\hat{\varepsilon}^2 - \varepsilon^2) \delta_t \tilde{w}_0 = O(M^{-1} \ln^m M). \quad (4.3)$$

Let  $M_j^{(l)} := \max_{(x,t) \in [0,1] \times [t_{j-1}, t_j]} |\frac{\partial^l}{\partial t^l} \tilde{w}_0|$ . Taylor series expansions yield  $|R_1| \leq C\varepsilon^2 \min\{k_j M_j^{(2)}, M_j^{(1)}\}$ ; similarly, we get  $|R_2| \leq C(\hat{\varepsilon}_j^2 - \varepsilon^2)k_j^{-1} M_j^{(0)}$ . Note that, by (3.16), here we have  $M_j^{(l)} \leq C\varepsilon^{-2l} e^{-\gamma^2 t_{j-1}/\varepsilon^2}$  as  $\frac{\partial^l}{\partial t^l} = \varepsilon^{-2l} \frac{\partial^l}{\partial \tau^l}$ . Therefore

$$|R_1| \leq C \min\left\{\frac{k_j}{\varepsilon^2}, 1\right\} e^{-\gamma^2 t_{j-1}/\varepsilon^2}, \quad |R_2| \leq C \max\{0, \hat{C} - \varepsilon^2 k_j^{-1}\} e^{-\gamma^2 t_{j-1}/\varepsilon^2}. \quad (4.4)$$

We shall show that (4.4) implies (4.3) for the Bakhvalov mesh (a) and the Shishkin mesh (b) separately.

(a) Case 1:  $\frac{j}{M} \leq \theta_0 - C_5 M^{-1}$  for some  $C_5 \geq 1$ . Then we have  $k_j \leq M^{-1} t'(\frac{j}{M}) = M^{-1} \frac{\varepsilon^2}{\gamma^2} [\frac{1}{2} - \frac{j}{M}]^{-1}$  and  $e^{-\gamma^2 t_{j-1}/\varepsilon^2} = [1 - 2\frac{(j-1)}{M}]$ , which imply

$$|R_1| \leq CM^{-1} \frac{1 - 2\frac{(j-1)}{M}}{\frac{1}{2} - \frac{j}{M}} = CM^{-1} 2 \left(1 + \frac{M^{-1}}{\frac{1}{2} - \frac{j}{M}}\right) \leq CM^{-1}.$$

Here we used  $\frac{1}{2} - \frac{j}{M} \geq C_5 M^{-1}$ , which follows from  $\theta_0 - \frac{j}{M} \geq C_5 M^{-1}$ . Furthermore, choosing  $C_5 = C_5(\hat{C})$  sufficiently large, we obtain  $\hat{C}k_j \leq \varepsilon^2$ , which yields  $R_2 = 0$ .

Case 2:  $\frac{j}{M} > \theta_0 - C_5 M^{-1} = \frac{1}{2} - C_4 \varepsilon^2 - C_5 M^{-1}$ . Now, a calculation shows that  $t_{j-1} = t(\frac{j-1}{M}) \geq \frac{\varepsilon^2}{\gamma^2} \ln(\frac{1}{2}[C_4 \varepsilon^2 + (C_5 + 1)M^{-1}]^{-1})$  and thus  $e^{-\gamma^2 t_{j-1}/\varepsilon^2} \leq C(\varepsilon^2 + M^{-1})$ . Combining this with  $\min\{\frac{k_j}{\varepsilon^2}, 1\} \leq C\varepsilon^{-2} \min\{M^{-1}, \varepsilon^2\}$  and the observation that  $\hat{C} - \varepsilon^2 k_j^{-1} > 0$  implies  $\varepsilon^2 \leq CM^{-1}$ , yields the bounds  $|R_1| \leq CM^{-1}$  and  $|R_2| \leq CM^{-1}$ , respectively.

(b) For  $j \leq M/2$  we have  $k_j = C\varepsilon^2 M^{-1} \ln M$ , which implies  $R_1 \leq CM^{-1} \ln M$  and  $R_2 = 0$  (as in this case  $k_j \leq \varepsilon^2/\hat{C}$  for sufficiently large  $M$ ). Otherwise, for  $j > M/2$ , we get  $e^{-\gamma^2 t_{j-1}/\varepsilon^2} \leq e^{-\gamma^2 \sigma_0/\varepsilon^2} = M^{-1}$  and thus  $|R_1| \leq CM^{-1}$  and  $|R_2| \leq CM^{-1}$ .  $\square$

## 4.2 Existence and accuracy, discrete upper and lower solutions

To establish existence of solutions of semilinear discrete equations (1.3) and (1.4), we invoke the theory of discrete upper and lower solutions outlined in the following result.

**PROPOSITION 4.2** Assume that on an arbitrary mesh  $\{(x_i, t_j)\}$  there exist discrete functions  $\alpha$  and  $\beta$  such that  $\alpha_{ij} \leq \beta_{ij}$  and

$$\hat{\mathcal{J}}^h \alpha_{ij} \leq 0 \leq \hat{\mathcal{J}}^h \beta_{ij}, \quad \alpha_{i,0} \leq \varphi(x_i) \leq \beta_{i,0}, \quad \alpha_{0,j} \leq g_0(t_j) \leq \beta_{0,j}, \quad \alpha_{N,j} \leq g_1(t_j) \leq \beta_{N,j},$$

where  $i = 1, \dots, N-1$ ,  $j = 1, \dots, M$ . Then problem (1.4) has a solution  $\hat{U}_{ij}$  such that  $\alpha_{ij} \leq \hat{U}_{ij} \leq \beta_{ij}$ .

*Proof.* The desired result is obtained imitating the proof of (Pao, 1985, Theorem 3.1) (where the case of  $\hat{\varepsilon}_j = \varepsilon = 1$  was considered). It is crucial in this argument that the discrete operator  $\hat{\mathcal{J}}^h + CI$  satisfies the discrete maximum principle, where  $I$  is the identity operator and  $C$  is an arbitrarily large but fixed positive constant. Alternatively, one can get the assertion of this proposition noting that the mapping  $\hat{\mathcal{J}}^h: \mathbb{R}^{(N+1)+(M+1)} \rightarrow \mathbb{R}^{(N+1)+(M+1)}$  is a Z-field; see Lorenz (1981); Kopteva & Stynes (2004).  $\square$

**REMARK 4.3** The functions  $\alpha$  and  $\beta$  of Proposition 4.2 are called ordered discrete lower and upper solutions (or sub- and super-solutions) of the discrete problem (1.4).

Now we are prepared to state existence and  $\varepsilon$ -uniform accuracy of our discrete solutions.




FIG. 2. Solution of test problem (5.1) for  $\varepsilon = 10^{-4}$  (left); a layer-adapted mesh with  $N = M = 16$  (right).

**THEOREM 4.4** Let the mesh  $\{(x_i, t_j)\}$  be either the Bakhvalov mesh of §4.1(a), or the Shishkin mesh of §4.1(b). Then for  $N$  and  $M$  sufficiently large, there exist solutions  $U_{ij}$  and  $\hat{U}_{ij}$  of discrete problems (1.3) and (1.4), respectively, such that for all  $i = 0, \dots, N$ ,  $j = 0, \dots, M$  we have

$$\begin{aligned} |U_{ij} - u(x_i, t_j)| &\leq C(N^{-2} \ln^{2m} N + M^{-1} \ln^m M), \\ |\hat{U}_{ij} - u(x_i, t_j)| &\leq C(N^{-2} \ln^{2m} N + M^{-1} \ln^m M), \end{aligned}$$

where  $m = 0$  for the Bakhvalov mesh (a) and  $m = 1$  for the Shishkin mesh (b).

*Proof.* As problem (1.3) is a particular case of problem (1.4), it suffices to prove the desired assertions only for  $\hat{U}_{ij}$ . Set  $\bar{p} = C_6(N^{-2} \ln^{2m} N + M^{-1} \ln^m M)$  and choose  $C_6$  sufficiently large so that, invoking Lemma 4.1, we get  $|\hat{\mathcal{T}}^h \beta - \mathcal{T} \beta| \leq C_0 \bar{p}/2$  for all  $|p| \leq p_0$ . In particular, this estimate holds for  $\beta(x, t; \pm \bar{p})$ , as for sufficiently large  $N$  and  $M$  we have  $\bar{p} \leq p_0$ . Furthermore, in view of (4.1), for sufficiently large  $N$  and  $M$ , we enjoy  $C_0 \bar{p} \gamma^2 - C_1(\varepsilon^2 + \bar{p}^2) \geq C_0 \bar{p}/2$ . Now, invoking Corollary 3.8 with  $p = \pm \bar{p}$ , we get  $\mathcal{T} \beta(x, t; -\bar{p}) \leq -C_0 \bar{p}/2$  and  $\mathcal{T} \beta(x, t; \bar{p}) \geq C_0 \bar{p}/2$ . These bounds immediately imply  $\hat{\mathcal{T}}^h \beta(x_i, t_j; -\bar{p}) \leq 0$  and  $\hat{\mathcal{T}}^h \beta(x_i, t_j; \bar{p}) \geq 0$ ; thus we obtained a discrete analogue of estimate (3.29a) in the proof of Theorem 3.9. Using (3.20) and (3.24), we now imitate the remaining part of this proof and conclude that  $\beta(x_i, t_j; -\bar{p})$  and  $\beta(x_i, t_j; \bar{p})$  are discrete lower and upper solutions. Furthermore, by Lemma 3.6 and Theorem 3.9, we have  $\beta(x_i, t_j; \pm \bar{p}) = u_{\text{as}} + O(\bar{p}) = u + O(\varepsilon^2 + \bar{p})$ . Finally, by Proposition 4.2, there exists  $\hat{U}_{ij}$  between  $\beta(x_i, t_j; -\bar{p})$  and  $\beta(x_i, t_j; \bar{p})$ ; therefore  $\hat{U}_{ij} = u(x_i, t_j) + O(\varepsilon^2 + \bar{p})$ , and recalling assumption (4.1), we get the desired estimate for  $\hat{U}_{ij}$ .  $\square$

## 5. Numerical results

Our model problem is (1.1) in the domain  $(x, t) \in [0, 1] \times [0, 2]$  with

$$f(x, t, u) := (2 - u)(u - u_1)u(u - u_2), \quad \text{where } u_1 := 1 - \frac{1}{2} \sin\left(\frac{\pi x}{2} - t\right), \quad u_2 := -(x^2 + \frac{1}{2}). \quad (5.1)$$

The corresponding reduced problem (1.2) has two stable solutions  $u_1$  and  $u_2$  and two unstable solutions 0 and 2. We use the boundary conditions  $g_0(t) = 0.6e^{-t} - 0.5$  and  $g_1(t) = 0.2e^{-t} - 0.1$ , and the initial condition  $\varphi(x) = 0.1$ . A calculation shows that the boundary conditions satisfy (A2) for both  $u_0 = u_1$





Table 2. Stabilized method (1.4) with  $\hat{C} = 4$  on the Bakhvalov mesh. Computational rates  $r$  in  $(N^{-1})^r$  (upper part) and maximum nodal errors (lower part).

$N$	$\varepsilon = 10^{-1}$	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-3}$	$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-5}$	$\varepsilon = 10^{-6}$	$\varepsilon = 10^{-7}$	$\varepsilon = 10^{-8}$	$\max_{\varepsilon}$
32	1.94	2.05	2.06	2.06	2.06	2.06	2.06	2.06	2.06
64	2.02	1.99	1.99	1.99	1.99	1.99	1.99	1.99	1.99
128	2.00	2.05	2.00	2.00	2.00	2.00	2.00	2.00	2.00
32	7.98e-3	1.97e-2	2.00e-2	2.01e-2	2.01e-2	2.01e-2	2.01e-2	2.01e-2	2.01e-2
64	2.08e-3	4.76e-3	4.80e-3	4.81e-3	4.81e-3	4.81e-3	4.81e-3	4.81e-3	4.81e-3
128	5.14e-4	1.19e-3	1.21e-3	1.21e-3	1.21e-3	1.21e-3	1.21e-3	1.21e-3	1.21e-3
256	1.28e-4	2.88e-4	3.01e-4	3.02e-4	3.02e-4	3.02e-4	3.02e-4	3.02e-4	3.02e-4

Table 3. Conventional method (1.3) on the Shishkin mesh. Computational rates  $r$  in  $(N^{-1} \ln(N/4))^r$  (upper part) and maximum nodal errors (lower part).

$N$	$\varepsilon = 10^{-1}$	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-3}$	$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-5}$	$\varepsilon = 10^{-6}$	$\varepsilon = 10^{-7}$	$\varepsilon = 10^{-8}$	$\max_{\varepsilon}$
32	3.34	2.45	2.45	2.45	2.45	2.45	2.45	2.45	2.45
64	2.96	2.04	2.04	2.04	2.04	2.04	2.04	2.04	2.04
128	2.71	2.00	2.01	2.01	2.01	2.01	2.01	2.01	2.00
32	9.37e-3	4.07e-2	4.07e-2	4.07e-2	4.07e-2	4.07e-2	4.07e-2	4.07e-2	4.07e-2
64	2.42e-3	1.51e-2	1.51e-2	1.51e-2	1.51e-2	1.51e-2	1.51e-2	1.51e-2	1.51e-2
128	6.03e-4	5.77e-3	5.77e-3	5.78e-3	5.78e-3	5.78e-3	5.78e-3	5.78e-3	5.78e-3
256	1.51e-4	2.07e-3	2.07e-3	2.07e-3	2.07e-3	2.07e-3	2.07e-3	2.07e-3	2.07e-3

Table 4. Stabilized method (1.4) with  $\hat{C} = 4$  on the Shishkin mesh. Computational rates  $r$  in  $(N^{-1} \ln(N/4))^r$  (upper part) and maximum nodal errors (lower part).

$N$	$\varepsilon = 10^{-1}$	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-3}$	$\varepsilon = 10^{-4}$	$\varepsilon = 10^{-5}$	$\varepsilon = 10^{-6}$	$\varepsilon = 10^{-7}$	$\varepsilon = 10^{-8}$	$\max_{\varepsilon}$
32	3.22	2.55	2.56	2.56	2.56	2.56	2.56	2.56	2.56
64	2.96	2.03	2.03	2.03	2.03	2.03	2.03	2.03	2.03
128	2.71	1.99	1.99	1.99	1.99	1.99	1.99	1.99	1.99
32	8.92e-3	4.13e-2	4.14e-2	4.14e-2	4.14e-2	4.14e-2	4.14e-2	4.14e-2	4.14e-2
64	2.42e-3	1.47e-2	1.47e-2	1.47e-2	1.47e-2	1.47e-2	1.47e-2	1.47e-2	1.47e-2
128	6.03e-4	5.67e-3	5.67e-3	5.67e-3	5.67e-3	5.67e-3	5.67e-3	5.67e-3	5.67e-3
256	1.51e-4	2.06e-3	2.05e-3	2.05e-3	2.05e-3	2.05e-3	2.05e-3	2.05e-3	2.06e-3

and  $M$ , a solution on an auxiliary mesh was used with  $2N$  and  $4M$  mesh intervals in the space and time directions, respectively).

Examining Tables 1–4, we conclude that the errors stabilize as  $\varepsilon$  approaches 0 and, furthermore, the convergence rates confirm the sharpness of the bounds of Theorem 4.4. Comparing the conventional method (1.3) and the stabilized method (1.4), we observe that although the errors of the stabilized method are slightly larger on the Bakhvalov mesh, on the considered layer-adapted meshes both the methods enjoy quite similar  $\varepsilon$ -uniform accuracy.

## 6. Conclusions

We have shown that the conventional implicit method (1.3) might produce incorrect and unstable computed solutions on uniform meshes; see Figures 1 and 3. Therefore we propose a stabilized method (1.4), which involves a constant parameter  $\hat{C} \geq 0$ . For this method, Proposition 1.2 prescribes a choice of  $\hat{C}$  that ensures uniqueness of the computed solution. Furthermore, our numerical results suggest that under this choice of  $\hat{C}$ , switching to the stabilized method cures the instability and yields qualitatively correct computed solutions on any mesh.

We theoretically investigated these two methods on layer-adapted meshes of Bakhvalov and Shishkin types and established their second-order convergence (with, in the case of the Shishkin mesh, a logarithmic factor) in the discrete maximum norm, uniformly in  $\varepsilon$  for  $\varepsilon \leq C(N^{-1} + M^{-1/2})$ ; see Theorem 4.4.

Although both the considered methods yield accurate computed solutions on layer-adapted meshes, we note that the conventional method (1.3) is unstable on certain meshes, which might be unacceptable, e.g., if a layer-adapted mesh is constructed adaptively, starting from an unsophisticated initial mesh. Therefore we advocate the stabilized method (1.4) over the conventional method (1.3).

## REFERENCES

- BAKHVALOV, N.S. (1969) Towards optimization of methods for solving boundary value problems in the presence of a boundary layer, *Zh. Vychisl. Mat. i Mat. Fiz.*, **9**, 841–859 (in Russian).
- FIFE, P.C. (1973) Semilinear elliptic boundary value problems with small parameters, *Arch. Rational Mech. Anal.*, **52**, 205–232.
- GRINDROD, P. (1991) *Patterns and Waves: the theory and applications of reaction-diffusion equations*. Oxford: Clarendon Press.
- KOPTEVA, N. (2007) Maximum norm error analysis of a 2d singularly perturbed semilinear reaction-diffusion problem, *Math. Comp.*, **76**, 631–646.
- KOPTEVA, N. & STYNES, M. (2004) Numerical analysis of a singularly perturbed nonlinear reaction-diffusion problem with multiple solutions, *Appl. Numer. Math.*, **51**, 273–288.
- LORENZ, J. (1981) Nonlinear singular perturbation problems and the Enquist-Osher scheme, *Mathematical Institute, Catholic University of Nijmegen*, Report 8115 (unpublished).
- MILLER, J.J.H., O’RIORDAN, E. & SHISHKIN, G.I. (1996) *Fitted Numerical Methods for Singular Perturbation Problems*. Singapore: World Scientific.
- MURRAY, J.D. (1993) *Mathematical Biology*. Berlin: Springer-Verlag.
- NEFEDOV, N.N. (1995) The method of differential inequalities for some classes of nonlinear singularly perturbed problems with internal layers, (*Russian*) *Differ. Uravn.*, **31**, 1142–1149; translation in *Differ. Equ.*, 1077–1085.
- PAO, C.V. (1985) Monotone iterative methods for finite difference system of reaction-diffusion equations, *Numer. Math.*, **46**, 571–586.
- PAO, C.V. (1992) *Nonlinear parabolic and elliptic equations*. New York: Plenum Press.
- PROTTER, M.H. & WEINBERGER, H.F. (1999) *Maximum principles in differential equations*. New York: Springer-Verlag.
- ROOS, H.-G., STYNES, M. & TOBISKA, L. (2008) *Robust Numerical Methods for Singularly Perturbed Differential Equations*. Berlin: Springer-Verlag.
- SHISHKIN, G.I. (1992) *Grid approximation of singularly perturbed elliptic and parabolic equations*. Ekaterinburg: Ur. O. Ran (in Russian).
- SUN, G. & STYNES, M. (1996) A uniformly convergent method for a singularly perturbed semilinear reaction-diffusion problem with multiple solutions, *Math. Comp.*, **65**, 1085–1109.
- VASIL’EVA, A.B., BUTUZOV, V.F. & KALACHEV, L.V. (1995) *The boundary function method for singular perturbation problems*. Philadelphia: SIAM.